

B<sup>1</sup>  
conceded

acetate phthalate (CAP), hydroxypropylmethyl cellulose phthalate, hydroxymethyl cellulose acetate succinate, methacrylic acid copolymers (e.g., Eudragit L30D-55 (trade name; produced by Rohm), Colicoat MAE30DP (trade name; produced by BASF), Polyquid PA30 (trade name; produced by San-yo Chemical)), carboxymethylethyl cellulose and shellac; sustained-release substrates such as methacrylic acid polymers (e.g., Eudragit NE30 D (trade name), Eudragit RL30D (trade name), Eudragit RS30D (trade name), etc.); water-soluble polymers; plasticizers such as triethyl citrate, polyethylene glycol, acetylated monoglycerides, triacetin and castor oil; and mixtures thereof. The above-mentioned "additive" includes, for example, water-soluble sugar alcohols (e.g., sorbitol, mannitol, maltitol, reduced starch saccharides, xylitol, reduced palatinose, erythritol, etc.), crystalline cellulose (e.g., Ceolas KG 801, Avicel PH 101, Avicel PH 102, Avicel PH 301, Avicel PH 302, Avicel RC-591 (crystalline cellulose carmellose sodium)), low-substituted hydroxypropyl cellulose (e.g., LH-22, LH-32, LH-23, LH-33 (Shin-Etsu Chemical) and mixtures thereof); binders, souring agents, bubbling agents, sweetening agents, flavorings, lubricants, coloring agents, stabilizers, excipients, disintegrants etc. are also used.

Please substitute the following paragraph for the fourth paragraph on page 14 of the specification.

Page 14, fourth paragraph (AMENDED)

B<sup>2</sup>

Lansoprazole (racemate) (34.2 g) was dissolved in 2-propanol (1,710 ml) and hexane (1,140 ml) containing triethylamine (0.2%) and fractionated by HPLC (column: CHIRALCEL OD 50 mm dia. X 500 mm, temperature: room temperature, mobile phase: hexane/2-propanol = 85/15, flow rate: 60 ml/min, detection wavelength: 285 nm, single injection: about 300 mg) to isolate the individual optical isomers. Fractions of an optical isomer of shorter retention time were combined and concentrated; the individual lots were combined and dissolved in ethanol